

Complete Summary

GUIDELINE TITLE

Tobacco control.

BIBLIOGRAPHIC SOURCE(S)

Michigan Quality Improvement Consortium. Tobacco control. Southfield (MI): Michigan Quality Improvement Consortium; 2005 Sep. 1 p.

GUIDELINE STATUS

This is the current release of the guideline.

This guideline updates a previous version: Michigan Quality Improvement Consortium. Tobacco control. Southfield (MI): Michigan Quality Improvement Consortium; 2003 Sep. 1 p.

COMPLETE SUMMARY CONTENT

SCOPE
 METHODOLOGY - including Rating Scheme and Cost Analysis
 RECOMMENDATIONS
 EVIDENCE SUPPORTING THE RECOMMENDATIONS
 BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS
 QUALIFYING STATEMENTS
 IMPLEMENTATION OF THE GUIDELINE
 INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT
 CATEGORIES
 IDENTIFYING INFORMATION AND AVAILABILITY
 DISCLAIMER

SCOPE

DISEASE/CONDITION(S)

Tobacco use

GUIDELINE CATEGORY

Evaluation
 Management
 Treatment

CLINICAL SPECIALTY

Family Practice
Internal Medicine

INTENDED USERS

Advanced Practice Nurses
Health Plans
Physician Assistants
Physicians

GUIDELINE OBJECTIVE(S)

- To achieve significant, measurable improvements in the management of tobacco control through the development and implementation of common evidence-based clinical practice guidelines
- To design concise guidelines that are focused on key management components of tobacco control to improve outcomes

TARGET POPULATION

- All patients 12 years of age and older (regardless of prior use status)
- All patients identified as current smokers/tobacco users

INTERVENTIONS AND PRACTICES CONSIDERED

Evaluation

Assessment of tobacco use status

Management/Treatment

1. Assessment of willingness to quit
2. Self-help material
3. Nicotine replacement therapy for adults
4. Withdrawal medication (e.g., sustained release bupropion) for adolescents and adults
5. Smoking cessation program
6. Follow-up contact

MAJOR OUTCOMES CONSIDERED

Not stated

METHODOLOGY

METHODS USED TO COLLECT/SELECT EVIDENCE

Searches of Electronic Databases

DESCRIPTION OF METHODS USED TO COLLECT/SELECT THE EVIDENCE

The Michigan Quality Improvement Consortium (MQIC) project leader conducts a search of current literature in support of the guideline topic. Computer database searches are used to identify published studies and existing protocols and/or clinical practice guidelines on the selected topic. A database such as MEDLINE and two to three other databases are used.

NUMBER OF SOURCE DOCUMENTS

Not stated

METHODS USED TO ASSESS THE QUALITY AND STRENGTH OF THE EVIDENCE

Weighting According to a Rating Scheme (Scheme Given)

RATING SCHEME FOR THE STRENGTH OF THE EVIDENCE

Levels of Evidence for the Most Significant Recommendation

- A. Randomized controlled trials
- B. Controlled trials, no randomization
- C. Observational studies
- D. Opinion of expert panel

METHODS USED TO ANALYZE THE EVIDENCE

Review

DESCRIPTION OF THE METHODS USED TO ANALYZE THE EVIDENCE

Not stated

METHODS USED TO FORMULATE THE RECOMMENDATIONS

Expert Consensus

DESCRIPTION OF METHODS USED TO FORMULATE THE RECOMMENDATIONS

Using the health plan guideline summaries and information obtained from the literature search, the Michigan Quality Improvement Consortium (MQIC) director and/or project leader prepare a draft guideline for review by the MQIC Medical Directors.

The draft guideline and health plan guideline summaries are distributed to the MQIC Medical Directors for review and discussion at their next committee meeting.

The review/revision cycle may be conducted over several meetings before consensus is reached. Each version of the draft guideline is distributed to the

MQIC Medical Directors, Measurement, and Implementation committee members for review and comments. All feedback received is distributed to the entire membership.

Once the MQIC Medical Directors achieve consensus on the draft guideline, it is considered approved for external distribution to practitioners with review and comments requested.

RATING SCHEME FOR THE STRENGTH OF THE RECOMMENDATIONS

Not applicable

COST ANALYSIS

A formal cost analysis was not performed and published cost analyses were not reviewed.

METHOD OF GUIDELINE VALIDATION

External Peer Review

DESCRIPTION OF METHOD OF GUIDELINE VALIDATION

Once the Michigan Quality Improvement Consortium (MQIC) Medical Directors achieve consensus on the draft guideline, it is considered approved for external distribution to practitioners with review and comments requested.

The MQIC director also forwards the approved guideline draft to presidents of the appropriate state medical specialty societies for their input. All feedback received from external reviews is presented for discussion at the next MQIC Medical Directors Committee meeting. In addition, physicians are invited to attend the committee meeting to present their comments.

RECOMMENDATIONS

MAJOR RECOMMENDATIONS

The level of evidence grades (A-D) are provided for the most significant recommendations and are defined at the end of the "Major Recommendations" field.

All Patients 12 Years of Age and Older (Regardless of Prior Use Status)

Identification of tobacco use status (never, former, current) and type (all forms, including smokeless tobacco, pipe, snuff, and cigars)

- Ask and record tobacco use status in the medical record and/or problem list [A].

Frequency

- At each outpatient visit and inpatient admission

All Patients Identified as Current Smokers/Tobacco Users

Intervention to promote cessation of tobacco use

- Advise to quit [A].
- Assess patient willingness to attempt to quit [C].
- Assist patients who are ready to quit by:
 - Establishing a quit date
 - Providing self-help materials
 - Offering nicotine replacement therapy (adults only) and/or withdrawal medications (e.g., sustained release bupropion) [A] (adolescents and adults)
 - Offering referral into smoking cessation program
- Arrange follow-up contact, either in person or by telephone [D]:
 - First week after quit date
 - First month after quit date

Frequency

- At each periodic health exam, more frequently at the discretion of the physician

Special Circumstances

- Pregnant Smokers: Due to the serious risks to the mother and fetus, pregnant smokers should be offered interventions such as referral to a smoking cessation program.
- Hospitalized Smokers: Clinicians should provide appropriate pharmacotherapy and counseling during hospitalization to reduce nicotine withdrawal symptoms and assist smokers in quitting.
- Smokers with Psychiatric Comorbidity: Nicotine withdrawal symptoms may exacerbate depression among patients with a prior history of affective disorder. Stopping smoking may affect the pharmacokinetics of certain psychiatric agents. Clinicians should monitor closely the actions or side effects of psychiatric medications in smokers/tobacco users who are attempting to quit.

Definitions:

Levels of Evidence for the Most Significant Recommendation

- A. Randomized controlled trials
- B. Controlled trials, no randomization
- C. Observational studies
- D. Opinion of expert panel

CLINICAL ALGORITHM(S)

None provided

EVIDENCE SUPPORTING THE RECOMMENDATIONS

TYPE OF EVIDENCE SUPPORTING THE RECOMMENDATIONS

The type of evidence is provided for the most significant recommendations (See "Major Recommendations" field).

This guideline is based on the 2000 U.S. Department of Health and Human Services guideline, Treating Tobacco Use and Dependence (www.surgeongeneral.gov/tobacco/).

BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS

POTENTIAL BENEFITS

Through a collaborative approach to developing and implementing common clinical practice guidelines and performance measures for tobacco control, Michigan health plans will achieve consistent delivery of evidence-based services and better health outcomes. This approach also will augment the practice environment for physicians by reducing the administrative burdens imposed by compliance with diverse health plan guidelines and associated requirements.

POTENTIAL HARMS

Not stated

QUALIFYING STATEMENTS

QUALIFYING STATEMENTS

This guideline lists core management steps. Individual patient considerations and advances in medical science may supersede or modify these recommendations.

IMPLEMENTATION OF THE GUIDELINE

DESCRIPTION OF IMPLEMENTATION STRATEGY

When consensus is reached on a final version of the guideline, a statewide mailing of the approved guideline is completed. The guideline is distributed to physicians in the following medical specialties:

- Family Practice
- General Practice
- Internal Medicine
- Other Specialists for which the guideline is applicable (e.g., endocrinologists, allergists, pediatricians, cardiologists)

INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES

IOM CARE NEED

Getting Better
Staying Healthy

IOM DOMAIN

Effectiveness
Patient-centeredness

IDENTIFYING INFORMATION AND AVAILABILITY

BIBLIOGRAPHIC SOURCE(S)

Michigan Quality Improvement Consortium. Tobacco control. Southfield (MI):
Michigan Quality Improvement Consortium; 2005 Sep. 1 p.

ADAPTATION

This guideline is based on the 2000 U.S. Department of Health and Human
Services guideline, Treating Tobacco Use and Dependence
(www.surgeongeneral.gov/tobacco/).

DATE RELEASED

2003 Sep (revised 2005 Sep)

GUIDELINE DEVELOPER(S)

Michigan Quality Improvement Consortium

SOURCE(S) OF FUNDING

Michigan Quality Improvement Consortium

GUIDELINE COMMITTEE

Michigan Quality Improvement Consortium Medical Director's Committee

COMPOSITION OF GROUP THAT AUTHORED THE GUIDELINE

Physician representatives from participating Michigan Quality Improvement
Consortium health plans, Michigan State Medical Society, Michigan Osteopathic
Association, Michigan Association of Health Plans, Michigan Department of
Community Health, and Michigan Peer Review Organization

FINANCIAL DISCLOSURES/CONFLICTS OF INTEREST

Not stated

GUIDELINE STATUS

This is the current release of the guideline.

This guideline updates a previous version: Michigan Quality Improvement Consortium. Tobacco control. Southfield (MI): Michigan Quality Improvement Consortium; 2003 Sep. 1 p.

GUIDELINE AVAILABILITY

Electronic copies: Available in Portable Document Format (PDF) from the [Michigan Quality Improvement Consortium Web site](#).

AVAILABILITY OF COMPANION DOCUMENTS

None available

PATIENT RESOURCES

None available

NGC STATUS

This NGC summary was completed by ECRI on April 14, 2004. The information was verified by the guideline developer on July 27, 2004. This NGC summary was updated by ECRI on November 28, 2005. The updated information was verified by the guideline developer on December 19, 2005.

COPYRIGHT STATEMENT

This NGC summary is based on the original guideline, which may be reproduced with the citation developed by the Michigan Quality Improvement Consortium.

DISCLAIMER

NGC DISCLAIMER

The National Guideline Clearinghouse™ (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC Inclusion Criteria which may be found at <http://www.guideline.gov/about/inclusion.aspx>.

NGC, AHRQ, and its contractor ECRI make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.

© 1998-2006 National Guideline Clearinghouse

Date Modified: 9/25/2006

